



Analysis of 2011 data

4329.0.00.006

Mortality of People Using

~~Mental Health Services~~
PSYCHIATRISTS! ←

and

~~Prescription Medications~~
PSYCHIATRIST'S KILLING DRUGS! ←

KEY FINDINGS :

- Persons who accessed mental health-related treatments accounted for 49.4% of all deaths in this period **75,858 deaths**

See the 'HORRIFIC' psychiatrist mass-killing death rates! ← **PRESS**

THIS GOVERNMENT REPORT STATES THAT 76,000 PEOPLE PER YEAR WERE 'KILLED' BY PSYCHIATRIC-DRUG-PUSHER PSYCHIATRISTS IN 2011; IT IS NOW OVER 80,000 !!

HEROIN DRUG-PUSHERS ONLY 'KILL' 750 AUSTRALIANS PER YEAR !!

PSYCHIATRIC-DRUG-PUSHER PSYCHIATRISTS 'KILL' 10,700 PERCENT 'MORE' !!

VICTORIAN PSYCHIATRIC-DRUG-PUSHER PSYCHIATRISTS 'FORCE' 700% MORE THAN USA !

AUSTRALIAN BUREAU OF STATISTICS

EMBARGO: 11.30AM (CANBERRA TIME) FRIDAY 08 SEPTEMBER 2017

Mortality of People Using Mental Health Services and Prescription Medications, Analysis of 2011 data



ARCHIVED ISSUE Released at 11:30 AM (CANBERRA TIME)
08/09/2017 **First Issue**

- [Downloads](#)
- [Explanatory Notes](#)
- [Related Information](#)
- [Past Releases](#)

Page tools: [Print Page](#) [Print All](#)

KEY FINDINGS

Characteristics of deaths of persons who accessed mental health-related treatments in 2011

Persons of all ages

- There were 3.2 million persons who accessed MBS and/or PBS subsidised mental health-related services or prescription medications ('mental health-related treatments') in 2011.
- **There were 153,451 deaths registered in Australia in the period 10 August 2011 to 27 September 2012.**
- **Persons who accessed mental health-related treatments accounted for 49.4% of all deaths in this period (75,858 deaths).**
- The standardised death rate for persons who accessed mental health-related treatments in 2011 was almost twice (1.9 times) that of the standardised death rate for the total Australian population (11.4 deaths per 1,000 population compared with 6.1 deaths per 1,000 population respectively).